



## TAF II p135/p105 Polyclonal Antibody

Cat No: HR1AP8446

For research use only

### Overview

Product Name	TAF II p135/p105 Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,ELISA
Species Reactivity	Human,Mouse
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	110kDa
GenelD?Human?	TAF4/TAF4B
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	TAF4; TAF2C; TAF2C1; TAF4A; TAFII130; TAFII135; Transcription initiation factor TFIID subunit 4; RNA polymerase II TBP-associated factor subunit C; TBP-associated factor 4; Transcription initiation fa
Background	TATA-box binding protein associated factor 4(TAF4) Homo sapiens Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the larger subunits of TFIID that has been shown to potentiate transcriptional activation by retinoic acid, thyroid hormone and