

MDC1 (phospho Ser513) Polyclonal Antibody

Cat No: HR1AP9192

For research use only

Overview

Product Name	MDC1 (phospho Ser513) Polyclonal Antibody
Source	Rabbit
Applications	IHC-p,ELISA
Species Reactivity	Human
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	kDa
GeneID?Human?	MDC1
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	MDC1; KIAA0170; NFB1; Mediator of DNA damage checkpoint protein 1; Nuclear factor with BRCT domains 1
Background	mediator of DNA damage checkpoint 1(MDC1) Homo sapiens The protein encoded by this gene contains an N-terminal forkhead domain, two BRCA1 C-terminal (BRCT) motifs and a central domain with 13 repetitions of an approximately 41-amino acid sequence. The encoded protein is required to activate the intra-S phase and G2/M phase cell cycle checkpoints in response to DNA damage. This nuclear protein interacts with phosphorylated histone H2AX near sites of DNA double-strand breaks through its BRCT motifs, and facilitates recruitment of the ATM kinase and meiotic recombination 11 protein complex to DNA damage foci. [provided by RefSeq, Jul 2008],