


COX1/Cyclooxygenase 1 Rabbit pAb

Cat No: HR1APEA023

For research use only

Overview

Product Name	COX1/Cyclooxygenase 1 Rabbit pAb
Source	Rabbit
Applications	WB, IHC
Species Reactivity	Human
Recommended Dilutions	WB 1:1,000 IHC 1:100
Immunogen	
Species	
Storage	
Isotype	
Clonality	Polyclonal
Concentration	1mg/ml
Observed band	70kDa
GeneID?Human?	5742
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	N/A
Background	Cyclooxygenase-1 (COX-1), also known as prostaglandin G/H synthase 1, prostaglandin-endoperoxide synthase 1 or prostaglandin H2 synthase 1, is an enzyme that in humans is encoded by the PTGS1 gene. There are two isoforms of COX encoded by distinct gene products: a constitutive COX-1 (this enzyme) and an inducible COX-2, which differ in their regulation of expression and tissue distribution. The expression of these two transcripts is differentially regulated by relevant cytokines and growth factors. A splice variant of COX-1 termed COX-3 was identified in the CNS of dogs, but does not result in a functional protein in humans. Two smaller COX-1-derived proteins (the partial COX-1 proteins PCOX-1A and PCOX-1B) have also been discovered, but their precise roles are yet to be described. Galectin 3 is one of the more extensively studied members of this family and is a 30 kDa protein. Due to a C-terminal carbohydrate binding site, Galectin 3 is capable of binding IgE and mammalian cell surfaces only when homodimerized or homooligomerized. Galectin 3 is normally distributed in epithelia of many organs, in various inflammatory cells, including macrophages, as well as dendritic cells and Kupffer cells. The expression of this lectin is up-regulated during inflammation, cell proliferation, cell differentiation and through trans-activation by viral proteins.