

TRAP220 Polyclonal Antibody

Cat No: HR1AP4751

For research use only

Overview

Product Name	TRAP220 Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,ELISA
Species Reactivity	Human,Mouse
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	170kDa
GeneID?Human?	MED1
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	MED1; ARC205; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; Mediator of RNA polymerase II transcription subunit 1; Activator-recruited cofactor 205 kDa component; ARC2
Background	mediator complex subunit 1(MED1) Homo sapiens The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. It also regulates p53-dependent apoptosis and it is essential for adipogenesis. This protein is known to have the ability to self-oligomerize. [provided by RefSeq, Jul 2008],