

## PDGFR? (5D1) Mouse mAb

Cat No: HR1AM2447

For research use only

## Overview

Product Name	PDGFR? (5D1) Mouse mAb
Source	Mouse
Applications	IHC
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	IHC 1:100-200
Immunogen	Al of the second
Species	Mouse
Storage	PBS with 0.02% sodium azide and 50% glycerol pH 7.4. Store at -20°C. Avoid repeated freeze-thaw cycles.
Isotype	lgG1
Clonality	Monoclonal
Concentration	1mg/ml
Observed band	180kDakDa
GeneID?Human?	5156
Human Swiss- Prot No.	
Cellular localization	
Alternative Names	N/A
Background	Platelet derived growth factor (PDGF) family proteins exist as several disulphide-bonded, dimeric isoforms (PDGF AA, PDGF AB, PDGF BB, PDGF CC, and PDGF DD) that bind in a specific pattern to two closely related receptor tyrosine kinases, PDGF receptor? (PDGFR?) and PDGF receptor? (PDGFR?) and PDGFR? and PDGFR? can each form heterodimers with EGFR, which is also activated by PDGF. Various cells differ in the total number of receptors present and in the receptor subunit composition, which may account for responsive differences among cell types to PDGF binding. Ligand binding induces receptor dimerization and autophosphorylation, followed by binding and activation of cytoplasmic SH2 domain-containing signal transduction molecules, such as GRB2, Src, GAP, PI3 kinase, PLC?, and NCK. A number of different signaling pathways are initiated by activated PDGF receptors and lead to control of cell growth, actin reorganization, migration, and differentiation.