

Endophilin I Polyclonal Antibody

Cat No: HR1AP8295

For research use only

Overview

Product Name	Endophilin I Polyclonal Antibody
Source	Rabbit
Applications	WB,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	39kDa
GeneID?Human?	SH3GL2
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	SH3GL2; CNSA2; SH3D2A; Endophilin-A1; EEN-B1; Endophilin-1; SH3 domain protein 2A; SH3 domain-containing GRB2-like protein 2
Background	domain:An N-terminal amphipathic helix, the BAR domain and a second amphipathic helix inserted into helix 1 of the BAR domain (N-BAR domain) induce membrane curvature and bind curved membranes. The BAR domain dimer forms a rigid crescent shaped bundle of helices with the pair of second amphipathic helices protruding towards the membrane-binding surface.,function:Implicated in synaptic vesicle endocytosis. May recruit other proteins to membranes with high curvature.,miscellaneous:HeLa cells expressing the N-BAR domain of SH3GL2 show tubulation of the plasma membrane. The N-BAR domain binds liposomes and induces formation of tubules from liposomes. The N-terminal amphipathic helix is required for liposome binding. The second amphipathic helix enhances liposome tubulation.,similarity:Belongs to the endophilin family.,similarity:Contains 1 BAR domain.,similarity:Contains 1 SH3 domain.,subcellular location:Concentrated in presynaptic nerve terminals in neurons.,subunit:Monomer; in cytoplasm. Homodimer; when associated with membranes (By similarity). Interacts with SYNJ1 and DNM1. Interacts with MAP4K3; the interaction appears to regulate MAP4K3-mediated JNK activation. Interacts with PDCD6IP.,tissue specificity:Brain, mostly in frontal cortex. Expressed at high level in fetal cerebellum.,