



Bag-4 Polyclonal Antibody

Cat No: HR1AP2856

For research use only

Overview

Product Name	Bag-4 Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	50kDa
GenID?Human?	BAG4
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	BAG4; SODD; BAG family molecular chaperone regulator 4; BAG-4; Bcl-2-associated athanogene 4; Silencer of death domains
Background	BCL2 associated athanogene 4(BAG4) Homo sapiens The protein encoded by this gene is a member of the BAG1-related protein family. BAG1 is an anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. This protein was found to be associated with the death domain of tumor necrosis factor receptor type 1 (TNF-R1) and death receptor-3 (DR3), and thereby negatively regulates downstream cell death signaling. The regulatory role of this protein in cell death was demonstrated in epithelial cells which undergo apoptosis while integrin mediated matrix contacts are lost. Alternatively spliced transcript variants encoding distinct