



Bag-3 Polyclonal Antibody

Cat No: HR1AP5385

For research use only

Overview

Product Name	Bag-3 Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,ELISA
Species Reactivity	Human
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	65kDa
GenID?Human?	BAG3
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	BAG3; BIS; BAG family molecular chaperone regulator 3; BAG-3; Bcl-2-associated athanogene 3; Bcl-2-binding protein Bis; Docking protein CAIR-1
Background	BCL2 associated athanogene 3(BAG3) Homo sapiens BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008],