

PKA α cat (phospho Thr198) Polyclonal Antibody

Cat No: HR1AP7851

For research use only

Overview

Product Name	PKA α cat (phospho Thr198) Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,IF,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	40kDa
GeneID?Human?	PRKACA/PRKACB
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	PRKACA; PKACA; cAMP-dependent protein kinase catalytic subunit alpha; PKA C-alpha; PRKACB; cAMP-dependent protein kinase catalytic subunit beta; PKA C-beta; PRKACG; cAMP-dependent protein kinase catal
Background	protein kinase cAMP-activated catalytic subunit alpha(PRKACA) Homo sapiens This gene encodes one of the catalytic subunits of protein kinase A, which exists as a tetrameric holoenzyme with two regulatory subunits and two catalytic subunits, in its inactive form. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. cAMP-dependent phosphorylation of proteins by protein kinase A is important to many cellular processes, including differentiation, proliferation, and apoptosis. Constitutive activation of this gene caused either by somatic mutations, or genomic duplications of regions that include this gene, have been associated with hyperplasias and adenomas of the adrenal cortex and are linked to corticotropin-independent Cushing's syndrome. Altern