

LIMK-2 (phospho Thr505) Polyclonal Antibody

Cat No: HR1AP2464

For research use only

Overview

Product Name	LIMK-2 (phospho Thr505) Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,IF,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	kDa
GeneID?Human?	LIMK2
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	LIMK2; LIM domain kinase 2; LIMK-2
Background	LIM domain kinase 2(LIMK2) Homo sapiens There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],