

## **Recombinant Mouse IL3**

Cat No:HR2R2168

For research use only

## Overview

Quantity	10 ?g
Gene Symbol	IL3
Gene ID	16187
Accession	P01586
Alternative Name	IL-3, Mast cell growth factor, MCGF, Multipotential colony-stimulating factor, Multi-CSF, HCGF, P-cell stimulation factor, Hematopoietic growth factor stimulation factor, Multi-CSF, HCGF, P-cell stimulation factor, Hematopoietic growth factor stimulation factor, Multi-CSF, HCGF, P-cell stimulation factor, Hematopoietic growth factor stimulation factor, Multi-CSF, HCGF, P-cell stimulation factor, Hematopoietic growth factor 
Species	Mouse
Source	E. coli
Description	IL3 is produced mainly by T cells following cell activation by antigens and mitogens, but also by keratinocytes, natural killer cells, mast cells, endothelial cells, and monocytes. The analysis of bacterial-derived recombinant IL3 shows that glycosylation is not required for the activity of IL3. IL-3 sequences are evolutionarily less well conserved. Human and murine IL3 show approximately 29% homology at the protein level while murine and rat IL3 show approximately 54% homology. IL3 receptors are expressed on macrophages, mast cells, eosinophils, megakaryocytes, basophils, bone marrow progenitor cells, and various myeloid leukemia cells. Binding of IL3 to its receptor causes specific phosphorylation of a 150 kDa membrane glycoprotein. Recombinant mouse IL3 is a monomer of 15 kDa.
Functions	The ED50 as determined by the dose-dependent stimulation of the proliferation of mouse M-NFS-60 cells was found to be in the <0.05 ng/mL
Formulation	Lyophilized from a 0.2 ?m filtered solution in PBS
Solubility	A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Appearance	Lyophilized Powder
Molecular Weight	15
Purity	>95% as determined by SDS-PAGE
Concentration	<1.0 EU/?g of recombinant protein as determined by the LAL method
Shipping Condition	Ambient Temperature
Storage Condition	The lyophilized protein is stable for at least one year from date of receipt at -70?C. Upon reconstitution, this cytokine can be stored in working aliquots at 2? - 8?C for one month, or at -20?C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles www.bioelsa.com info@bioelsa.com

info@bioelsa.com