

Recombinant Human FGF1 Cat No:HR2R1382

For research use only

Overview

Quantity	1.0 ?g
Gene Symbol	FGF1
Gene ID	9158
Accession	P05230
Alternative Name	aFGF, FGF-1, Heparin-binding growth factor 1, HBGF-1, Endothelial cell growth factor, ECGF-beta, FGFA, FGF1
Species	Human
Source	E. coli
Description	aFGF and bFGF display 55% homology which is limited, however, to a few functional domains. The structure of aFGF, revealed by X-ray crystallography, shows a similar folding structure as IL-1. Because aFGF binds to the same receptor as bFGF, aFGF displays more or less the same spectrum of activities. It is, however, generally less active than bFGF in similar assays. FGF receptors are encoded by a gene family consisting of at least four receptor tyrosine kinases that transduce signals important in a variety of developmental and physiological processes related to cell growth and differentiation.
Functions	The ED(50) as determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF receptors is ? 10 ng/mL, corresponding to a specific activity of ? 1.0 x 10^5 units/mg.
Formulation	Recombinant acidic FGF was lyophilized from a 0.2 ?m filtered solution in 2.5% glycine, 0.5% sucrose, 0.01% Tween80, 5 mM Glutamic acid, pH 4.5.
Solubility	A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Appearance	Lyophilized Powder
Molecular Weight	16
Purity	>95% as determined by SDS-PAGE
Concentration	<1.0 EU/?g of recombinant protein as determined by the LAL method.
Shipping Condition	Ambient Temperature
Storage Condition	The lyophilized protein is stable for at least one year from date of receipt at -70?C. Upon reconstitution, this cytokine can be stored in working aliquots at 2? - 8?C for one month, or at -20?C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.