

## Recombinant Human FGR (N-GST tag) Cat No:HR2R1424

For research use only

## Overview

Quantity	10 ?g
Gene Symbol	FGR
Gene ID	N/A
Accession	NM_005248
Alternative Name	SRC2; c-fgr; c-src2; FLJ43153; MGC75096; p55c-fgr; p58c-fgr
Species	Human
Source	
Description	FGR is a protooncogene that is a unique member of the tyrosine kinase gene family. Certain lymphomas (but not sarcomas or carcinomas) express FGR-related messenger RNA. This transcript is detected in Burkitt's lymphoma cell lines naturally infected with Epstein-Barr virus (EBV), but not in EBV-negative Burkitt's lymphoma cells . FGR expression is limited to normal peripheral blood granulocytes, monocytes, and alveolar macrophages, all of which contain 50 to 100 copies of c-fgr mRNA per cell .?
Functions	The specific activity of FGR was determined to be 289 nmol /min/mg as per activity assay protocol.
Formulation	50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.
Solubility	N/A
Appearance	Liquid
Molecular Weight	86
Purity	70% - 90%
Concentration	
Shipping Condition	Dry Ice
Storage Condition	Store product at ?70?C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.