

Recombinant Human IL10

Cat No:HR2R1535

For research use only

Overview

| | |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Quantity | 25 ?g |
| Gene Symbol | IL10 |
| Gene ID | 3586 |
| Accession | P22301 |
| Alternative Name | IL-10, Cytokine synthesis inhibitory factor, CSIF, B-TCGF, TGIF Recombinant Human Interleukin-10 (IL10) |
| Species | Human |
| Source | CHO cells |
| Description | <p>IL-10 is produced by mouse Th2 cells following their stimulation by lectins. The main source for B cell derived IL-10 in mice are Ly1 B cells that express CD5 (Ly1) and CD11. In humans IL-10 is produced by activated CD8(+) peripheral blood T cells, by T helper CD4(+) T cell clones after both antigen-specific and polyclonal activation, by B cell lymphomas, and by monocytes following cell activation by bacterial lipopolysaccharides and mast cells. IL-10 is a homodimeric protein with subunits having a length of 160 amino acids. Human Interleukin-10 shows 73% amino acid homology with mouse Interleukin-10. The murine IL-10 receptor has been cloned. This receptor is a protein of approximately 110 kDa that binds murine IL-10 specifically. This receptor is structurally related to receptors for IFN. The CD nomenclature for this receptor is CDw210. IL-10 inhibits the synthesis of a number of cytokines such as IFN-gamma, IL-2 and TNF-beta in Th1 T helper subpopulations of T cells but not of Th2 T helper cells. This activity is antagonized by IL-4. The inhibitory effect on IFN-gamma production is indirect and appears to be the result of a suppression of IL-12 synthesis by accessory cells. In the human system, IL-10 is produced by, and down-regulates the function of Th1 and Th2 cells.</p> |
| Functions | The ED(50) determined by dose-dependent co-stimulation (with murine IL-4) of MC/9 cells. The ED(50) was found to be ? 2.0 ng/mL, corresponding to a specific activity of ? 4.0 x 10 ⁵ units/mg. |
| Formulation | Recombinant Interleukin-10 was lyophilized from a 0.2 ?m filtered PBS solution pH 7.0. |
| Solubility | A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers. |
| Appearance | Lyophilized Powder |
| Molecular Weight | 19 |
| Purity | >95% as determined by SDS-PAGE |
| Concentration | <1.0 EU/?g of recombinant protein as determined by the LAL method. |
| Shipping Condition | Ambient Temperature |

Storage
Condition

The lyophilized protein is stable for at least one year from date of receipt at -70°C. Upon reconstitution, this cytokine can be stored in working aliquots at 2° - 8°C for one month, or at -20°C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.

