

## **Recombinant Human PDGFB** Cat No:HR2R1834

For research use only

## Overview

Quantity	1.0 ?g
Gene Symbol	PDGFB
Gene ID	5155
Accession	P01127
Alternative Name	PDGF subunit B, PDGF-2, Platelet-derived growth factor B chain, Platelet-derived growth factor beta polypeptide, Proto-oncogene c-Sis, NN=Becaplermin br/>Recombinant Human Platelet-Derived Growth Factor Subunit B (PDGFB)
Species	Human
Source	
Description	The PDGF family is comprised of five different disulphide-linked dimers of four different polypeptide chains: A, B, C and D (PDGF-AA, PDGF-BB, PDGF-AB, PDGF-CC and PDGF-DD). Synthesized mainly by megakaryocytes, PDGFs are stored in the alpha granules of platelets from which they are released following platelet activation. Functioning as an autocrine and paracrine growth factor, PDGFs are involved in a number of biological processes that include but not limited to hyperplasia, chemotaxis, embryonic neuron development, wound healing and respiratory tubule epithelial cell development. Aberrant expression of PDGFs is observed with vascular proliferative diseases such as atherosclerosis. PDGFs regulate the synthesis of their own receptor and also influence the expression of membrane receptors for IL1, EGF, 5-Hydroxytryptamine, LDL and transferrin. Recombinant human PDGFB is a disulfide-linked homodimer of two B chains.
Functions	The ED50 as determined by the dose-dependent proliferation of NIH 3T3 cells is <0.5ng/ml
Formulation	Lyophilized from a 0.2 ?m filtered solution in PBS
Solubility	A quick spin of the vial followed by reconstitution in sterile distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Appearance	Lyophilized Powder
Molecular Weight	12.3
Purity	>95% as determined by SDS-PAGE
Concentration	<1.0 EU/?g of recombinant protein as determined by the LAL method.
Shipping Condition	Ambient Temperature

Storage	
Condition	

The lyophilized protein is stable for at least one year from date of receipt at -70?C. Upon reconstitution, this cytokine can be stored in working aliquots at 2? - 8?C for one month, or at -20?C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.

