

Recombinant Human QIK (N-GST tag)

Cat No:HR2R1908

For research use only

Overview

Quantity	10 ?g
Gene Symbol	QIK
Gene ID	N/A
Accession	NM_015191
Alternative Name	SNF1LK2, SIK2, KIAA0781, LOH11CR1I, DKFZp434K1115
Species	Human
Source	Insect cells
Description	QIK is a serine/threonine protein kinase that contains an N-terminal kinase domain, a central domain with ubiquitin-associate motif, and a C-terminal PKA phosphorylation site. QIK can phosphorylate IRS1 and overexpression of QIK in adipocyte elevates the phosphorylation of IRS1. The QIK-mediated phosphorylation of IRS1 may modulate the efficiency of insulin signal transduction and could be responsible for insulin resistance associated with diabetes. Insulin disrupts TORC2 activity by induction of QIK which then stimulates the phosphorylation and cytoplasmic translocation of TORC2. Phosphorylated TORC2 is subsequently degraded by the 26S proteasome.
Functions	The specific activity of QIK was determined to be 112nmol /min/mg as per activity assay protocol.
Formulation	50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol.
Solubility	N/A
Appearance	Liquid
Molecular Weight	150
Purity	70% - 90%
Concentration	
Shipping Condition	Dry Ice
Storage Condition	Store product at ?70?C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.