

Recombinant Human TOPK (N-GST tag)

Cat No:HR2R2029

For research use only

Overview

Quantity	10 ?g
Gene Symbol	ТОРК
Gene ID	N/A
Accession	NM_018492
Alternative Name	PBK, SPK, Nori-3, FLJ14385
Species	Human
Source	Insect cells
Description	TOPK is a MAPK kinase that phosphorylates p38 MAPK and is activated in a cell-cycle-dependent manner in neuronal progenitor cells in vitro . Expression of TOPK is detected in male germ line progenitor cells, activated T-cells, and a variety of lymphomas and leukemias. In vitro studies have shown that activated TOPK phosphorylated p38MAPK but not JNK or ERK. TOPK activation requires phosphorylation by both the M-phase CDK1/CyclinB kinase complex and another unknown kinase, possibly RafC or RafA. TOPK may play an important role in linking extracellular signals to an intracellular state, possibly allowing extracellular influence on the cell-cycle-related processes of proliferation or differentiation .
Functions	The specific activity of TOPK was determined to be 7.5 nmol /min/mg as per activity assay protocol.
Formulation	50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.
Solubility	N/A
Appearance	Liquid
Molecular Weight	68
Purity	70% - 90%
Concentration	
Shipping Condition	Dry Ice
Storage Condition	Store product at ?70?C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.