

Recombinant Human TXK (N-GST tag) Cat No:HR2R2047

For research use only

Overview

Quantity	10 ?g
Gene Symbol	ТХК
Gene ID	N/A
Accession	NM_003328
Alternative Name	RLK; TKL; BTKL; PTK4; PSCTK5; MGC22473
Species	Human
Source	
Description	TXK or RLK is a member of the TEC family of non-receptor tyrosine kinases. TXK is expressed in T-cells and is an important component of signaling pathways downstream of lymphocyte antigen receptor. TXK is phosphorylated in response to T-cell receptor stimulation and can be activated by phosphorylation by Src family kinases. However, TXK is phosphorylated independent of PI-3K activity . Excessive TXK protein expression is seen in patients with Behcet's disease. Over production of TXK leads to increased Th1 cell function that is involved in the pathogenesis of Behcet?s disease .?
Functions	The specific activity of TXK was determined to be 18 nmol /min/mg as per activity assay protocol.
Formulation	50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.
Solubility	N/A
Appearance	Liquid
Molecular Weight	53
Purity	70% - 90%
Concentration	
Shipping Condition	Dry Ice
Storage Condition	Store product at ?70?C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.